

## References

- Alvarez, M. P., Samayoa-Mendez, A. X., Naglak, M. C., Yuschak, J. V., & Murayama, K. M. (2015). Risk factors for postoperative unplanned intubation: Analysis of a national database. *American Surgeon, 81*(8), 820-825.
- American College of Surgeons, NSQIP Operations Manual, Chapter 4, ACS NSQIP Variables and definitions, 2016. Retrieve from: <https://www.facs.org/quality-programs/acs-nsqip>
- Balas, M. C., Vasilevskis, E. E., Olsen, K. M., Schmid, K. K., Shostrom, V., Cohen, M. Z., . . . Burke, W. J. (2014). Effectiveness and safety of the awakening and breathing coordination, delirium monitoring/management, and early exercise/mobility bundle. *Critical Care Medicine, 42*(5), 1024-1036.
- Barr, J., Fraser, G., Puntillo, K., Ely, E., Gelinas, C., Dasta, J., . . . Skrobik, Y., (2013). Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Critical Care Medicine, 41*(1), 263-306.
- Brown, C.V., Daigle, J.B., Foulkrod, K.H., Brouillette, B., Clark, A., Czysz, C., . . . Cooper, H. (2011). Risk factors associated with early reintubation in trauma patients: A prospective observational study. *Journal of Trauma-Injury Infection & Critical Care, 71*(1), 37-41.
- Brueckmann, B., Villa-Uribe, J., Bateman, B.T., Grosse-Sundrup, M., Hess, D.R., Schlett, C.L., & Eikermann, M. (2013). Development and validation of a score for prediction of postoperative respiratory complications. *Anesthesiology, 118*(6), 1276-1285.
- Frutos-Vivar F., Ferguson N.D., Esteban A., Epstein S.K., Arabi Y., Apezteguia C, . . . Anzueto A. (2006). Risk factors for extubation failure in patients following a successful spontaneous breathing trial. *Chest, 130*(6), 1664-1671.

Healey, C., Osler, T. M., Rogers, F. B., Healey, M. A., Glance, L. G., Kilgo, P. D., . . . Meredith, J. W. (2003). Improving the glasgow coma scale score: Motor score alone is a better predictor. *Journal of Trauma-Injury Infection & Critical Care*, 54(4), 671-678.

Hua, M., Brady, J. E., & Li, G. (2012). A scoring system to predict unplanned intubation in patients having undergone major surgical procedures. *Anesthesia & Analgesia*, 115(1), 88-94.

Institute of Medicine: *To err is human*. Retrieved from

<https://iom.nationalacademies.org/~media/Files/Report%20Files/1999/To-Err-is-Human/To%20Err%20is%20Human%201999%20%20report%20brief.pdf>

Kiekkas, P., Aretha, D., Panteli, E., Baltopoulos, G.I., & Filos, K.S. (2013). Unplanned extubation in critically ill adults: Clinical review. *Nursing in Critical Care*, 18(3), 123-134.

McCaffrey, J., Farrell, C., Whiting, P., Dan, A., Bagshaw, S., Delaney, A., (2009). Corticosteroids to prevent extubation failure: a systematic review and meta-analysis, *Intensive Care Medicine*, 35:977-986.

Menon, N., Joffe, A.M., Deem, S., Yanez, N.D., Grabinsky, Dagal, A., . . . Treggiari, M.M. (2012). Occurrence and complications of tracheal reintubation in critically ill adults. *Respiratory Care*, 57(10), 1555-1563.

Miu, T., Joffe, A.M., Yanez, N.D., Khandelwal, N., Dagal, A., Deem, S., & Treggiari, M., (2014). Predictors of reintubation in critically ill patients. *Respiratory Care*, 59(2), 178-185.

- Mokhlesi, B., Tulaimat, A., Gluckman, T.J., Wang, Y., Evans, A.T., & Corbridge, T.C. (2007). Predicting extubation failure after successful completion of a spontaneous breathing trial. *Respiratory Care*, 52(12), 1710-1717.
- Politano, A., Riccio, L., Lake, D., Rusin, C., Guin, L., Josef, C., Clark, M., ... Calland, J., (2013). Predicting the Need for Urgent Intubation in a Surgical/Trauma Intensive Care Unit, *Surgery*, 154(5): 1110-1116.
- Pronovost, P. J., Goeschel, C.A., Marsteller, J.A., Sexton J.B., Pham J.C., & Berenholtz S.M. (2009). Framework for patient safety research and improvement. *Circulation*, 119(2), 330-337.
- Ramachandran, S. K., Nafiu, O. O., Ghaferi, A., Tremper, K. K., Shanks, A., & Kheterpal, S. (2011). Independent predictors and outcomes of unanticipated early postoperative tracheal intubation after nonemergent, noncardiac surgery. *Anesthesiology*, 115(1), 44-53.
- Rujirojindakul, P., Geater, A.F., McNeil, E.B., Vasinanukorn, P., Prathep, S., Asim, W., & Naklongdee, J., (2012). Risk factors for reintubation in the post-anaesthetic care unit: A case-control study. *British Journal of Anaesthesia*, 109(4), 636-642.
- Sessler, C. N., Gosnell, M. S., Grap, M. J., Brophy, G. M., O'Neal, P. V., Keane, K. A., . . . Elswick, R. K. (2002). The richmond agitation-sedation scale: Validity and reliability in adult intensive care unit patients. *American Journal of Respiratory & Critical Care Medicine*, 166(10), 1338-1344.
- Snyder, C. W., Patel, R. D., Roberson, E. P., & Hawn, M. T. (2009). Unplanned intubation after surgery: Risk factors, prognosis, and medical emergency team effects. *American Surgeon*, 75(9), 834-838.

Thille, A.W., Boissier, F., Ben Ghezala, H., Razazi, K., Mekontso-Dessap, A., & Brun-Buisson, C. (2014). Risk factors for and prediction by caregivers of extubation failure in ICU patients: A prospective study\*. *Critical Care Medicine*, 43(3), 613-620.